

Investigating the potential of *Cymodocea nodosa* (Ucria) Ascherson as a coastal carbon sink coupling marine habitat cartographies and *in situ* non-destructive sampling

Arnaud Abadie*^{1,2}, Jonathan Richir*^{3,1}, Riccardo Pieraccini⁴, Branko Velimirov⁵, Michèle Leduc², Gilles Lepoint¹, Sylvie Gobert¹

* First co-authors

¹ Laboratory of Oceanology, MARE Centre, University of Liege, B6C, 4000 Liege, Sart Tilman, Belgium

² STARESO SAS, Pointe Revellata, BP 33, F-20260 Calvi, France

³ Numerical Ecology of Aquatic Systems, Complexys Institute, University of Mons, Pentagone 3D08, 6, Avenue du Champ de Mars, 7000 Mons, Belgium

⁴ DISTAV, University of Genova, Corso Europa 26, 16132 Genova, Italy

⁵ Medical University of Vienna, Center for Pathobiochemistry and Genetics, Währingerstr.10, 1090 Vienna, Austria

Framework

CONTEXT: Seagrass meadows are **major carbon sinks**, a main focus being made on climax species like *Posidonia oceanica* (L.) Delile while other species remained little studied.

AIM: To investigate the carbon stocking capacity of *Cymodocea nodosa* (Ucria) Ascherson leaves (Fig. 1a) a **pioneer species** with a rapid turnover and an expected high **carbon stocking capacity**.

APPROACH:

- Estimation of the area covered by *C. nodosa* by **mapping marine habitats**.
- Sampling its leaves with **non-destructive** method for **biometry** as well as **carbon and nitrogen analysis**.

STUDY SITE: Calvi Bay (Fig. 1b), Corsica Island (Mediterranean Sea, France).




Figure 1: a) *C. nodosa* meadow in Corsica (France) (photo: A. Abadie); b) Location of the study site.

Marine habitats

In Calvi Bay, *C. nodosa* meadows cover an area of **0.498 km²** (Fig. 4). This value seems **underestimated** due to the difficulty to identify on side scan sonar images sparse meadows settled on sand.




Figure 4: Map of marine habitats in Calvi Bay (Corsica, France). *C. nodosa* meadows correspond to blue polygons. Red circles show the sampling sites. Modified from Velimirov et al. (2016).

Carbon and nitrogen stocks

The intermediate depth range (11-20 m) stores more than **60 % of the total carbon** stored by *C. nodosa* in Summer and **more than 80 %** in Winter (Tab. 2). The greater storage capacity of this bathymetrical zone can be explained by its **lower exposure** to water movement, however with **enough light** to sustain the plant growth. The nitrogen storage capacity follows the same pattern, with a low amount of nitrogen stored in *C. nodosa* leaf biomass in Winter when leaves are **exported to other marine systems** by storms.

Table 2: Area covered by *C. nodosa* in Calvi Bay and the amount of carbon (C) and nitrogen (N) contained in its leaves in Winter and Summer according to the bathymetrical range.

Depth range	Area (m ²)	C Winter (kg)	N Winter (kg)	C Summer (kg)	N Summer (kg)
0 - 10 m	57 210	40,1	2,2	969,9	39,9
11 - 20 m	427 774	212,6	11,5	1 701,5	81,8
21 - 30 m	13 459	2,6	0,2	43,6	2,4
Total	49 8443	255,3	13,9	2 715,0	124,2

Works cited

- Agostini S, Pergent G, Marchand B (2003) Growth and primary production of *Cymodocea nodosa* in a coastal lagoon. Aquat Bot 76:185-193
- de los Santos CB, Vicencio-Ramsey B, Lepoint G, Remy F, Bouma TJ, Gobert S (2016) Ontogenic variation and effect of collection procedure on leaf biomechanical properties of Mediterranean seagrass *Posidonia oceanica* (L.) Delile. Marine Ecology
- Pieraccini R (2015) Temporal and spatial variation of *Cymodocea nodosa* in the Calvi bay; first approach. Undergraduate Program of Environmental Science, University of Genova, Genova, Italy
- Velimirov B, Lejeune P, Kirschner A, Jousseume M, Abadie A, Pête DC, Dauby P, Richir J, Gobert S (2016) Estimating carbon fluxes in a *Posidonia oceanica* system: Paradox of the bacterial carbon demand. Estuarine, Coastal and Shelf Science 171:23-34

Methods

MAPPING

Georeferenced data obtained with high resolution side scan sonar images (Fig. 2), aerial orthophotographs, bathymetry and **ground truths** by scuba diving.

Cartographies of marine habitats are produced via **Computer-aided design (CAD)**. The map raster is then vectorised and computed through a **Geographical Information System (GIS)** to calculate the **area** covered by *C. nodosa*.




Figure 2: Side scan sonar image of a *C. nodosa* meadow in Calvi Bay.

LEAF SAMPLING, BIOMETRY AND ELEMENTAL CONTENTS

Shoot density and **non-destructive sampling** (de los Santos et al. 2016, Fig. 3a) were measured in Winter and Summer 2015 by cutting leaves (Fig. 3b-c) at **several depths** (5, 11, 15 and 23 m) at **different locations** within the bay (Calvi Beach, Alga Bay and Oscelluccia Bay). Leaf length and epiphyte biomass are measured in laboratory. **Carbon and nitrogen contents** are measured with an elemental analyser (Fig. 3d).




Figure 3: a) Measure of the shoot density; b) and c) *C. nodosa* leaf cutting at 20 m depth (photos: M. Leduc); d) Elemental analyzer for the measure of carbon and nitrogen contents (photos: R. Pieraccini).

Morphological features

C. nodosa shows the strongest seasonal variations at the **lowest depths** (Tab. 1) in Alga and Oscelluccia. The quantity of carbon and nitrogen stored in its leaves **decreases with depth** (Fig. 5) with a maximum storage at 5 m depth.

Table 1: Mean (\pm SD) shoot density, leaf and epiphyte dry biomass of *C. nodosa* at all sites in Winter and Summer 2015. Modified from Pieraccini (2016).

SITE	Depth (m)	Leaf density (leaf.m ⁻²)		Leaf biomass (gDW.m ⁻²)		Epiphyte biomass (gDW.shoot ⁻¹ .m ⁻²)	
		Winter	Summer	Winter	Summer	Winter	Summer
Alga	5	346 (\pm 75)	438 (\pm 163)	5,8 (\pm 1,3)	43,58 (\pm 22,4)	1,594	22,096
Oscelluccia	11	258 (\pm 85)	374 (\pm 57)	5,2 (\pm 1,7)	29,391 (\pm 11,9)	2,199	11,47
Calvi Beach	11	120 (\pm 42)	92 (\pm 43)	2,5 (\pm 0,9)	5,77 (\pm 2,6)	2,082	4,32
Calvi Beach	15	-	120 (\pm 61)	-	10,07 (\pm 5,8)	-	13,856
Calvi Beach	20	220 (\pm 163)	146 (\pm 111)	2,6 (\pm 1,9)	7,70 (\pm 3,1)	6,765	42,197
Calvi Beach	23	-	142 (\pm 41)	-	8,24 (\pm 2,6)	-	16,575




Figure 5: Mean carbon and nitrogen content of *C. nodosa* leaves at different depths in Winter (blue) and Summer (green).

C. nodosa roles

IN CALVI BAY

When compared with *P. oceanica* meadows and the 900 tons of carbon stored in their leaves in Summer (Velimirov et al. 2016), the capacity of *C. nodosa* meadows to be a major carbon sink appears negligible. However, their capacity to act through the export of their leaves as **a source of nitrogen in Winter** for other nearby systems must not be denied.

AROUND CORSICA

C. nodosa meadow shoot densities in Calvi Bay are very low compared to other Corsican areas like Urbino lagoon where a carbon storage capacity of 843.7 gC m⁻² per year was reported at 1 m depth (Agostini et al. 2003). Moreover, **the surface they cover in lagoons and coastal areas appears to be largely underestimated** according to the last mapping campaign made around Corsica in 2015.